

LA-UR-19-20054

Approved for public release; distribution is unlimited.

Title: Quantum Resources for Information Processing

Author(s): Girolami, Davide

Intended for: Talk for research visit at Purdue Univ.

Issued: 2019-01-08

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.


Quantum Resources for Information Processing

Davide Girolami


West Lafayette, 17 January 2019



How quantum differs from classical?



Quantum Resources



Goal: Characterizing Quantum Resources

? Theoretical Quantification: $f_R(\rho)$ being monotone under free operations

? Demonstration of supraclassical performance: $f_R(\rho)$ is figure of merit in a task

? Experimental Detection:

$$f_R(\rho) = \langle O_{\text{exp}} \rangle, O_{\text{exp}} = O_{\text{exp}}^\dagger$$


(Some) Technical Objectives

I Evaluating the power of quantum devices (Th)

II Evaluating the power of quantum devices (Exp)

III (Future) Quantum Resources for Artificial Intelligence

I Evaluating the power of quantum devices (Th)



- System \mathcal{S} , described by ρ_N
- How to quantify correlations of order $2 \leq k \leq N$ in an N particle system ?

Covariances? No thanks

$$\langle X_1 X_2 \dots X_N \rangle_{\rho_N} - \prod_i \langle X_i \rangle_{\rho_N}$$


Don't detect **classical** correlations **X**

Can be created by local operations **X**

Can be created by fine graining **X**

D. Kaszlikowski, A. Sen, U. Sen, V. Vedral, and A. Winter, PRL 101, 070502 (2008); Z. Walczak, Comment; D. Kaszlikowski, *et al.*, Reply. Z. Walczak, PLA 374, 3999 (2010), C. H. Bennett, *et al.*, PRA 83, 012312 (2011)

Correlation hierarchy



Relative entropy of genuine multipartite correlations

- Correlations higher than k : distance to P_k

$$S^{k \rightarrow N}(\rho_N) := \min_{\sigma \in P_k} S(\rho_N || \sigma)$$

- Genuine k -partite correlations:

$$S^k(\rho_N) := S^{k-1 \rightarrow N}(\rho_N) - S^{k \rightarrow N}(\rho_N)$$

A Complexity measure

- ! Equally correlated states can have very different structure
- ? Single index classifying multipartite classical and quantum states

Weaving := weighted sum of genuine multipartite correlations


$$W_S(\rho_N) = \sum_{k=2}^N \omega_k S^k(\rho_N)$$

D. Girolami, T. Tufarelli, and C. Susa, Phys. Rev. Lett. 119, 140505 (2017)


Test, $\omega_k = k - 1$

State	\mathbf{W}_S
$[(00\rangle\langle 00 + 11\rangle\langle 11)/2]^{\otimes N/2}$	$N/2$
$[\frac{(00\rangle + 11\rangle)}{\sqrt{2}}]^{\otimes N/2}$	N
$(\sum_{i=1}^d ii\rangle / \sqrt{d})^{\otimes N/2}$	$N \log d$
$(0\rangle^{\otimes N} + 1\rangle^{\otimes N})/\sqrt{2}$	$\sim N \log N$
$\sum_i \mathcal{P}_i(0\rangle^{\otimes N/2} \otimes 1\rangle^{\otimes N/2})/\sqrt{\binom{N}{N/2}}$	$\sim N^2$

II Evaluating the power of quantum devices (Exp)




Usefulness = Superlinear Speed



- ★ $\text{Speed}(\Pi_i U_t^i \psi_N U_t^{i\dagger}) \geq f_{\text{linear}}(N) \Rightarrow \psi_N$ is entangled
- ★ $O(4^N)$ measurements required to reconstruct system state


Evaluating Speed with limited resources




- ★ Two system copies and $O(3^n)$ measurements are sufficient to evaluate the speed function, even if system state and dynamics are unknown

EXP scheme for n-qubit systems

$$V_{1,2,\dots,n} = \bigotimes_i V_i, \quad V_i = I_2 - 2 |\phi_-^i\rangle \langle \phi_-^i|$$




C. M. Alves and D. Jaksch, PRL 93, 110501 (2004), H. Jeong *et al.*, J. Opt. Soc. Am. B 31, 3057 (2014)




Devices: IBM Q

- <https://www.research.ibm.com/ibm-q/>
- 5-qubit, 16-qubit machines
- Superconducting qubits initialized in ground state at 15ml K, gate error= 10^{-3}
- remote access via Composer and Qiskit

Entanglement detection in ibmqx4





BUT constraints... split into two experiments



A comparison

$$\star \rho = p |\phi^+\rangle \langle \phi^+| + (1-p) |\phi^-\rangle \langle \phi^-|, p \in [0, 1], U_t = e^{-i\sigma_x t}, t = \pi/6$$




Left: ibmqx4; Right: optical setup, C. Zhang et al. PRA 96, 042327 (2017)

Original Scheme: D. Girolami, PRL 113, 170401 (2014)


Multipartite Entanglement detection

- ★ A state ψ_N is k -partite entangled iff it is not a tensor product of states describing $\leq k$ -subsystems
- ★ E.g. $\psi_3 \neq \psi_2 \otimes \psi_1 \Rightarrow \psi_3$ displays 3-partite entanglement
- ★ $\text{Speed}(\sum_i U_t^i \psi_N U_t^{i\dagger}) \geq ([N/k]k + (N - [N/k]k)^2)/4$,
 $k \geq 2 \Rightarrow \psi_N$ is at least k -partite entangled

Multipartite Entanglement detection




BUT constraints... split into three experiments




Detection of speed-up due to tripartite and bipartite Entanglement

- ★ $|GHZ\rangle = 1/\sqrt{2}(|000\rangle + |111\rangle)$, $U_t = e^{-i\sigma_x t}, e^{-i\sigma_z t}$, $t = \pi/6$
- ★ $\text{Speed}() \geq .75$ certifies bipartite entanglement, $\text{Speed}() \geq 1.25$ certifies tripartite entanglement
- ★ $\text{SpeedX}(\Pi_i U_t^i |GHZ\rangle) = 0.61 \pm 0.16$ **X**
- ★ $\text{SpeedZ}(\Pi_i U_t^i |GHZ\rangle) = 1.89 \pm 0.20$. **V**


Bonus: Bound to state engineering




III Quantum Resources for AI



How quantum systems think?



How quantum systems think?



Quantum Decision Making

- Quantize classical strategies (Markov decision processes, dynamic programming)
- Attack by quantizing the Bellman equation(s) and performance comparing
- Deliberation as a quantum walk, faster decision (smarter?)
- Testable in today quantum machines, e.g. IBM chips (D-Wave?)

Summary

- Quantum Information Processing as clever use of quantum resources
- Foundational and practical value
- A great record of successes, but also future applications

The End

Email: davegirolami@gmail.com

Website: <https://sites.google.com/site/davegirolami/>